

Cited in Biological Abstracts; Elsevier BIOBASE/Current Awareness in Biological Sciences; Current Contents / Life Sciences; EMBASE/Excerpta Medica; Index Medicus; Informaticus; Pascal M

CONTENTS

Research Papers

- Growth response of breast epithelial cells to estrogen is influenced by EGF
P. Briand, B.K. Lundholt, J. Skouv, A.E. Lykkesfeldt (Denmark) 1
- Activation of the insulin-like growth factor binding protein-1 promoter by progesterone receptor in decidualized human endometrial stromal cells
J. Gao, J. Mazella, A. Suwanichkul, D.R. Powell, L. Tseng (USA) 11
- Discordant expression of insulin-like growth factors and their receptor messenger ribonucleic acids in endometrial carcinomas relative to normal endometrium
R.N. Roy, A.H. Gerulath, A. Cecutti, B.R. Bhavnani (Canada) 19
- Retinoid and androgen regulation of cell growth, epidermal growth factor and retinoic acid receptors in normal and carcinoma rat prostate cells
F. Richter, H.F.S. Huang, M.-t. Li, D. Danielpour, S.-L. Wang, R.J. Irwin, Jr. (USA) 29
- Participation of protein kinase C α in 1,25-dihydroxy-vitamin D₃ regulation of chick myoblast proliferation and differentiation
D.A. Capiati, M.T. Téllez-Iñón, R.L. Boland (Argentina) 39
- Growth hormone and insulin-like growth factor-I enhance β -glucuronidase gene activation by androgen in mouse kidney
G.L. Niermann, G.L. Watson (USA) 47
- Stimulation of specific binding of [³H]-progesterone to bovine luteal cell-surface membranes: specificity of digitonin
G.S. Menzies, K. Howland, M.T. Rae, T.A. Bramley (UK) 57
- The HDQVH-motif in domain E of the estradiol receptor α is responsible for zinc-binding and zinc-induced hormone release
A. Humeny, D. Bökenkamp, H.H. Thole (Germany) 71
- Lack of responsiveness to TGF- β 1 in a thyroid carcinoma cell line with functional type I and type II TGF- β receptors and Smad proteins, suggests a novel mechanism for TGF- β insensitivity in carcinoma cells
N.-E. Heldin, D. Bergström, A. Hermansson, A. Bergenstråhle, A. Nakao, B. Westermarck, P. ten Dijke (Sweden) 79
- The first exon of the human *sc* gene contains an androgen responsive unit and an interferon regulatory factor element
A. Haelens, G. Verrijdt, E. Schoenmakers, P. Alen, B. Peeters, W. Rombauts, F. Claessens (Belgium) 91
- Nitric oxide inhibits human aldosteronogenesis without guanylyl cyclase stimulation
E.L. Kreklau, E.J. Carlson, J.G. Drewett (USA) 103
- Osteogenic protein-1 increases gene expression of vascular endothelial growth factor in primary cultures of fetal rat calvaria cells
L.-C.C. Yeh, J.C. Lee (USA) 113

(Contents continued on inside back cover)

CONTENTS
Direct

This journal is part of **ContentsDirect**, the *free* alerting service which sends tables of contents by e-mail for Elsevier Science books and journals. You can register for **ContentsDirect** online at: www.elsevier.nl/locate/contentsdirect



0303-7207(19990720)153:1:2;1-6

08040

MOLECULAR AND CELLULAR ENDOCRINOLOGY

Publication information: *Molecular and Cellular Endocrinology* (ISSN 0303-7207). For 1999, volumes 148–158 are scheduled for publication. Subscription prices are available upon request from the Publisher or from the Regional Sales Office nearest you or from this journal's website (<http://www.elsevier.nl/locate/mce>). Further information is available on this journal and other Elsevier Science products through Elsevier's website: (<http://www.elsevier.nl>). Subscriptions are accepted on a prepaid basis only and are entered on a calendar year basis. Issues are sent by standard mail (surface within Europe, air delivery outside Europe). Priority rates are available upon request. Claims for missing issues should be made within six months of the date of dispatch.

Orders, claims and product enquiries: please contact the Customer Support Department at the Regional Sales Office nearest you:

New York: Elsevier Science, P.O. Box 945, New York, NY 10159-0945, USA; Tel. (+1)212-633-3730 [Toll free number for North American customers: 1-888-4ES-4ES-INFO (437-4636)]; Fax (+1)212-633-3680; E-mail: usinfo-f@elsevier.com

Amsterdam: Elsevier Science, P.O. Box 211, 1000 AE Amsterdam, The Netherlands; Tel. (+31)20-4853757; Fax (+31)20-4853432; E-mail: nlinfo-f@elsevier.nl

Tokyo: Elsevier Science, 9-15 Higashi-Azabu 1-chome, Minato-ku, Tokyo 106-0044, Japan; Tel. (+81)3-5561-5033; Fax (+81)3-5561-5047; E-mail: info@elsevier.co.jp

Singapore: Elsevier Science, No. 1 Temasek Avenue, #17-01 Millenia Tower, Singapore 039192, Tel. (+65)434-3727; Fax (+65)337-2230; E-mail: asiainfo@elsevier.com.sg

Rio de Janeiro: Elsevier Science, Rua Sete de Setembro 111/16 Andar, 20050-002 Centro, Rio de Janeiro - RJ, Brazil; Tel. (+55)21-5095-340; Fax (+55)21-507-1991; E-mail: elsevier@campus.com.br [Note (Latin america): for orders, claims and help desk information, please contact the Regional Sales Office in New York as listed above]

Advertising information: Advertising orders and enquiries may be sent to: **Europe and ROW:** Rachel Gresle-Farthing, Elsevier Science Ltd., Advertising Department, The Boulevard, Langford Lane, Kidlington, Oxford OX5 1GB, UK; Tel.: +44 1865 843565; Fax: +44 1865 843976; E-mail: r.gresle-farthing@elsevier.co.uk. **USA and Canada:** Elsevier Science Inc., Mr. Tino DeCarlo, 655 Avenue of the Americas, New York, NY 10010-5107, USA; Tel.: +1 212 633 3815; Fax: +1 212 633 3820; E-mail: t.decarlo@elsevier.com. **Japan:** Elsevier Science K.K., Advertising Department, 1-9-15 Higashi-Azabu 1-chome, Minato-ku, Tokyo 106, Japan; Tel.: +81 3 55615033; Fax: +81 3 55615047.

If you need information about the journal, you can find all details (including full "Notes to Authors") on the World Wide Web: access under <http://www.elsevier.nl/locate/mce> or <http://www.elsevier.com/locate/mce>

(Contents continued from outside back cover)

Transplantation of primary bovine adrenocortical cells into <i>scid</i> mice M. Thomas, P.J. Hornsby (USA)	125
Co-ordinate expression of activin A and its type I receptor mRNAs during phorbol ester-induced differentiation of human K562 erythroleukemia cells K. Hildén, T. Tuuri, M. Erämaa, O. Ritvos (Finland)	137
Unbound rather than total concentration and saturation rather than unsaturation determine the potency of fatty acids on insulin secretion C. Warnotte, M. Nenquin, J.-C. Henquin (Belgium)	147
GLUT2 and glucokinase expression is coordinately regulated by sulfonylurea O. Porzio, L.N.J.L. Marlier, M. Federici, M.L. Hribal, R. Magnaterra, D. Lauro, A. Fusco, G. Sesti, P. Borboni (Italy)	155
Natural variants of the β isoform of the human glucocorticoid receptor do not alter sensitivity to glucocorticoids P. de Lange, J.W. Koper, A.O. Brinkmann, F.H. de Jong, S.W.J. Lamberts (The Netherlands)	163
Cloning of zebrafish activin type IIB receptor (ActRIIB) cDNA and mRNA expression of ActRIIB in embryos and adult tissues R.R. Garg, L. Bally-Cuif, S.E. Lee, Z. Gong, X. Ni, C.L. Hew, C. Peng (Canada, Germany, Singapore)	169
Molecular mechanism for cooperation between Sp1 and steroidogenic factor-1 (SF-1) to regulate bovine CYP11A gene expression Z. Liu, E. R. Simpson (USA)	183
Calendar	197
Announcement	
Notes to Authors	

